Populations are dynamic entities. Populations consist all of the species living within a specific area, and populations fluctuate based on a number of factors: seasonal and yearly changes in the environment, natural disasters such as forest fires and volcanic eruptions, and competition for resources between and within species. The statistical study of population dynamics, demography, uses a series of mathematical tools to investigate how populations respond to changes in their biotic and abiotic environments. Many of these tools were originally designed to study human populations. The term "demographics" is commonly used when discussing humans, all living populations can be studied using this approach.

# Population Size and Density

The study of any population usually begins by determining how many individuals of a particular species exist, and how closely associated they are with each other. Within a particular habitat, a population can be characterized by its population size, the total number of individuals, and its population density, the number of individuals within a specific area or volume. Population size and density are the two main characteristics used to describe and understand populations. For example, populations with more individuals may be more stable than smaller populations based on their genetic variability, and thus their potential to adapt to the environment. Alternatively, a member of a population with low population density (more spread out in the habitat), might have more difficulty finding a mate to reproduce compared to a population of higher density.


## **Species Distribution (Dispersion)**

In addition to measuring simple density, further information about a population can be obtained by looking at the distribution of the individuals. Species dispersion patterns (or distribution patterns) show the spatial relationship between members of a population within a

habitat at a particular point in time. In other words, they show whether members of the species live close together or far apart, and what patterns are evident when they are spaced apart.

Individuals in a population can be equally spaced apart, dispersed randomly with no predictable pattern, or clustered in groups. These are known as uniform, random, and clumped dispersion patterns, respectively. Uniform dispersion is observed in plants that secrete substances inhibiting the growth of nearby individuals (such as the release of toxic chemicals by the sage plant *Salvia leucophylla*, a phenomenon called allelopathy) and in animals like the penguin that maintain a defined territory. An example of random dispersion occurs with dandelion and other plants that have wind-dispersed seeds that germinate wherever they happen to fall in a favorable environment. A clumped dispersion may be seen in plants that drop their seeds straight to the ground, such as oak trees, or in animals that live in groups (schools of fish or herds of elephants). Clumped dispersions may also be a function of habitat heterogeneity. Thus, the dispersion of the individuals within a population provides more information about how they interact with each other than does a simple density measurement. Just as lower density species might have more difficulty finding a mate, solitary species with a random distribution might have a similar difficulty when compared to social species clumped together in groups.

Species may have uniform, random, or clumped distribution. Territorial birds such as penguins tend to have uniform distribution. Plants such as dandelions with wind-dispersed seeds tend to be randomly distributed. Animals such as elephants that travel in groups exhibit clumped distribution.



## **Demography**

While population size and density describe a population at one particular point in time, scientists must use demography to study the dynamics of a population. Demography is the statistical study of population changes over time: birth rates, death rates, and life expectancies. Each of these measures, especially birth rates, may be affected by the population characteristics. For example, a large population size results in a higher birth rate because more potentially reproductive individuals are present. In contrast, a large population size can also result in a higher death rate because of competition, disease, and the accumulation of waste. Similarly, a higher population density or a clumped dispersion pattern results in more potential reproductive encounters between individuals, which can increase birth rate. Lastly, a female-biased sex ratio (the ratio of males to females) or age structure (the proportion of population members at specific age ranges) composed of many individuals of reproductive age can increase birth rates.

In addition, the demographic characteristics of a population can influence how the population grows or declines over time. If birth and death rates are equal, the population remains stable. However, the population size will increase if birth rates exceed death rates; the population will decrease if birth rates are less than death rates. Life expectancy is another important factor; the length of time individuals remain in the population impacts local

resources, reproduction, and the overall health of the population. These demographic characteristics are often displayed in the form of a life table.

#### Life Tables

Life tables provide important information about the life history of an organism. Life tables divide the population into age groups and often sexes, and show how long a member of that group is likely to live. They are modeled after actuarial tables used by the insurance industry for estimating human life expectancy. Life tables may include the probability of individuals dying before their next birthday (i.e., their mortality rate), the percentage of surviving individuals dying at a particular age interval, and their life expectancy at each interval. An example of a life table is shown in (Figure) from a study of Dall mountain sheep, a species native to northwestern North America. Notice that the population is divided into age intervals (column A). The mortality rate (per 1000), shown in column D, is based on the number of individuals dying during the age interval (column B) divided by the number of individuals surviving at the beginning of the interval (Column C), multiplied by 1000.

mortality rate = number of individuals dyingnumber of individuals surviving x 1000mortality rat e = number of individuals dyingnumber of individuals surviving x 1000

For example, between ages three and four, 12 individuals die out of the 776 that were remaining from the original 1000 sheep. This number is then multiplied by 1000 to get the mortality rate per thousand.

mortality rate =  $12776 \times 1000 \approx 15.5$ mortality rate =  $12776 \times 1000 \approx 15.5$ 

As can be seen from the mortality rate data (column D), a high death rate occurred when the sheep were between 6 and 12 months old, and then increased even more from 8 to 12 years old, after which there were few survivors. The data indicate that if a sheep in this

# Demography-Population Size and Density, Life Tables, Dispersion, Survivorship

population were to survive to age one, it could be expected to live another 7.7 years on average, as shown by the life expectancy numbers in column E.

This life table of *Ovis dalli* shows the number of deaths, number of survivors, mortality rate, and life expectancy at each age interval for the Dall mountain sheep.

# Life Table of Dall Mountain Sheep<sup>1</sup>

Age interval (years)(A)	Number dying in age interval out of 1000 born (B)	Number surviving at beginning of age interval out of 1000 born (C)	per 1000 alive at	Life expectancy or mean lifetime remaining to those attaining age interval (E)
0-0.5	54	1000	54.0	7.06
0.5-1	145	946	153.3	_
1-2	12	801	15.0	7.7
2-3	13	789	16.5	6.8
3-4	12	776	15.5	5.9
4-5	30	764	39.3	5.0
5-6	46	734	62.7	4.2
6-7	48	688	69.8	3.4
7-8	69	640	107.8	2.6
8-9	132	571	231.2	1.9
9-10	187	439	426.0	1.3

This life table of *Ovis dalli* shows the number of deaths, number of survivors, mortality rate, and life expectancy at each age interval for the Dall mountain sheep.


# Life Table of Dall Mountain Sheep<sup>1</sup>

Age interval (years)(A)	Number dying in age interval out of 1000 born (B)	Number surviving at beginning of age interval out of 1000 born (C)	per 1000 alive at	Life expectancy or mean lifetime remaining to those attaining age interval (E)
10-11	156	252	619.0	0.9
11-12	90	96	937.5	0.6
12-13	3	6	500.0	1.2
13-14	3	3	1000	0.7

# Survivorship Curves

Another tool used by population ecologists is a survivorship curve, which is a graph of the number of individuals surviving at each age interval plotted versus time (usually with data compiled from a life table). These curves allow us to compare the life histories of different populations ((Figure)). Humans and most primates exhibit a Type I survivorship curve because a high percentage of offspring survive their early and middle years—death occurs predominantly in older individuals. These types of species usually have small numbers of offspring at one time, and they give a high amount of parental care to them to ensure their survival. Birds are an example of an intermediate or Type II survivorship curve because birds die more or less equally

at each age interval. These organisms also may have relatively few offspring and provide significant parental care. Trees, marine invertebrates, and most fishes exhibit a Type III survivorship curve because very few of these organisms survive their younger years; however, those that make it to an old age are more likely to survive for a relatively long period of time. Organisms in this category usually have a very large number of offspring, but once they are born, little parental care is provided. Thus these offspring are "on their own" and vulnerable to predation, but their sheer numbers assure the survival of enough individuals to perpetuate the species.



Survivorship curves show the distribution of individuals in a population according to age. Humans and most mammals have a Type I survivorship curve because death primarily occurs in the older years. Birds have a Type II survivorship curve, as death at any age is equally probable. Trees have a Type III survivorship curve because very few survive the younger years, but after a certain age, individuals are much more likely to survive.

## **Natality**

Natality is the ability of individuals of a population to produce new individuals. Natality rate is equivalent to the birth rate in the terminology of human population study

(Demography). Maximum (some time called absolute or physiological) natality is the theoretical maximum production of new individuals under ideal environmental conditions (i.e. no ecological limiting factors) and is a constant for a given population. Ecological or realized natality refers to population increase under an actual or specific environmental condition. It is not a constant for a population but may vary with the size and age composition of the population and the physical environmental conditions. Natality is generally expressed as a rate determined by dividing the number of new individuals produced by time (absolute or crude natality rate) or as the number of new individuals per unit of time per unit of population (the specific natality rate).

### Mortality

Natality refers to death of individuals in the population. Mortality is equivalent to death rate in human demography. Like natality, mortality may be expressed as the number of individuals dying in a given period (death per time). Ecological or realized mortality is the loss of individuals, under a given environmental condition. Like ecological natality, it is not a constant but varies with population and environmental conditions. A theoretical minimum mortality, a constant for a population, represents the loss under ideal or non limiting conditions.

### Age structure

It is obvious that individuals in a population will be of different age groups. Relative numbers of young and old individuals in a population will significantly influence the behavior of a population such as natality and mortality. Thye proportion of individuals in each age group is called age structure of that population. Age distribution influences both natality and motality of the population. The ratio of the various age groups in a population determines the current reproductive status of the population. The age structure of any population can be classified in to three categories i.e. pre-reproductive, reproductive and post-reproductive ages. The relative duration of these age groups in proportion to the life span varies greatly with different organisms. In man the three ages are relatively equal in length.

Reference: Mary Ann Clark, Jung Choi and Matthew Douglas, Biology